AD-A034 728 EDGEWOOD ARSENAL ABERDEEN PROVING GROUND MD
'IN VITRO' INHIBITION OF MICROSOMAL CALCIUM ATPASE FROM EGGSHEL--ETC(U)
DEC 76 G J KOLAJA, D E HINTON

UNCLASSIFIED

EB-TR-76097

NL

END
DATE
FILMED
3:14-77
NTIS


U.S. DEPARTMENT OF COMMERCE National Technical Information Service

AD-A034 728

'IN VITRO' INHIBITION OF MICROSOMAL CALCIUM ATPASE FROM EGGSHELL GLAND OF MALLARD DUCK

EDGEWOOD ARSENAL, ABERDEEN PROVING GROUND, MARYLAND

DECEMBER 1976



EDGEWOOD ARSENAL TECHNICAL REPORT

EB-TR-76097

IN VITRO INHIBITION OF MICROSOMAL CALCIUM ATPASE FROM EGGSHELL GLAND OF MALLARD DUCK

by


Gerald J. Kolaja, D.V.M., CPT, VC David E. Hinton, Ph.D.

Biomedical Laboratory

December 1976



DEPARTMENT OF THE ARMY
Headquarters, Edgewood Arsenal
Aberdeen Proving Ground, Maryland 21010



Approved for public release; distribution unlimited

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
. REPORT NUMBER	2. GOVT ACCESSION NO.	. 3. RECIPIENT'S CATALOG NUMBER
EP TD 74007		
EB-TR-76097 TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
		Technical Report
IN VITRO INHIBITION OF MICROSOM	IAL CALCIUM ATPASE	February 1976 - March 1976
FROM EGGSHELL GLAND OF MALLA	RD DUCK A	6. PERFORMING ORG. REPORT NUMBER
	***	6. PERFORMING ORG. REPORT NUMBER
AUTHOR(a)		S. CONTRACT OR GRANT NUMBER(*)
Gerald J. Kolaja, D.V.M., CPT VC David E. Hinton, Ph.D.	grand a story on and the course of the story of the course	
PERFORMING ORGANIZATION NAME AND ADD	DRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Commander, Edgewood Arsenal		Anen a work out i nombero
Attn: SAREA-BL-VP		EPA
Aberdeen Proving Ground, MD 21010		Add the control of th
. CONTROLLING OFFICE NAME AND ADDRESS	5	12. REPORT DATE
Commander, Edgewood Arsenal		December 1976
Attn: SAREA-TS-R		13. NUMBER OF PAGES
Aberdeen Proving Ground, MD 21010		10
4. MONITORING AGENCY NAME & ADDRESS(II	different from Controlling Office)	15. SECURITY CI.ASS. (of this report)
		UNCLASSIFIED
4		154. DECLASSIFICATION/DOWNGRADING SCHEDULE
		NA
Copies available from: National Technical 7. DISTRIBUTION STATEMENT (of the abetract of		
. SUPPLEMENTARY NOTES		
. KEY WORDS (Continue on reverse side if neces	eary and identify by block number;)
Ca ATPase		
DDT		
Microsomes		The
. ABSTRACT (Cantinue as reverse elde H naces	and identify by block number)	
		ATPase by DDT in an in vitro microson
test system. Furthermore, this inhibition		
to those of other investigators who show	ved Ca ATPase inhibition by	DDE in shell gland homogenate. Previ
studies have shown DDT-induced eggshe	ell thinning to be due to dec	creased calcium content of eggshells. Sin
calcium ATPase has been shown to be as	ssociated with calcium trans	port in eagshell sland, the inhibition of
enzyme in vitro offers a possible explanat	tion for DDT-induced eggshell	Atlanta
cital in the oriers a possible explanat	TOU TOU DO I MIGUICE CERTICIE	thinmix.

PREFACE

The work described in this report was authorized under Project/Task No. EPA. This work was started in February 1976 and completed in March 1976.

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals" as promulgated by the Committee on Revision of the Guide for Laboratory Animals Facilities and Care of the Institute of Laboratory Animal Resources, National Research Council.

The use of trade names in this report does not constitute an official endorsement or approval of the use of such commercial hardware or software. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, Edgewood Arsenal, Attn: SAREA-TS-R, Aberdeen Proving Ground, Maryland 21010; however DDC and the National Technical Information Service are authorized to reproduce the document for US Government purposes.

CONTENTS

		Page
I.	INTRODUCTION	5
II.	MATERIALS AND METHODS	5
III.	RESULTS	5
IV.	DISCUSSION	7
	LITERATURE CITED	8
	DISTRIBUTION LIST	9
	LIST OF FIGURES	
Figure		
1	In Vitro Ca ATPase Inhibition by DDT	6
2	Lineweaver-Burke Plot of Ca ATPase Inhibiton by DDT	6

DUC	White Section Buff Section
Unankorn Jestificat	350
DISTRIBUT	IGN/AVAILABILITY OGUES ATAIL DOS IT SPECIAL
*	I S COINE

IN VITRO INHIBITION OF MICROSOMAL CALCIUM ATPASE FROM EGGSHELL GLAND OF MALLARD DUCK

I. INTRODUCTION

In a previous paper we reported the effects of 2,2-bis(p-chlorophenyl)-1,1,1-trichloroethane (DDT) on morphology of eggshell gland of the mallard duck (Anas platy-rhynchos) (Kolaja and Hinton 1976). In that study, edema of the eggshell gland mucosa accompanied eggshell thinning. We postulated that the edema could have arisen as the result of an ionic imbalance. Calcium adenosine triphosphatase (ATPase) is the enzyme which is thought to be responsible for the transport of one ion, calcium, across epithelium of the eggshell gland (Pike and Alvarado 1975). Both in vivo and in vitro inhibition of Ca ATPase by the DDT metabolite 2,2-bis-(chlorophenyl)-1,1-dichloroethylene (DDE) has been demonstrated in homogenate of freeze-dried mucosal preparation from shell gland of Pekin ducks (Miller, et al., 1976). The primary purpose of this study was to determine the inhibitory effect of DDT on Ca ATPase isolated from microsomal preparations of duck eggshell glands and to characterize the type of inhibition by Lineweaver/Burke plots of Michaelis Menten analyses.

II. MATERIALS AND METHODS

Mature mallard hens, in egg production, were killed by cervical dislocation and the eggshell glands were rapidly excised and placed in ice-cold 0.25 M sucrose. All subsequent procedures were done in an ice bath or at 4°C. The mucosa of the eggshell gland was removed by scraping with a clean razor blade and 1 gm was weighed and diluted up to 10 ml in 0.25 M sucrose. The sucrose-shell gland mixture was homogenized at 500 rpm in a Potter-Elvehjem tissue homogenizer. The homogenate was sedimented for 15 minutes at 18,000 X g max in a Beckman L-2 ultracentrifuge. The above supernatant was recentrifuged at 98,000 X g max for 1 hr. The resulting pellet was resuspended to 3 ml in 0.25 M sucrose, and stored at -60°C until the time of assay. The Ca ATPase activity was determined by the method of Rorive and Kleinzeller (1974).4 Twenty microliters of DDT (Pfaltz and Bauer, Flushing, New York), dissolved in acetone were added to the reaction mixture and brought to a final concentration of 10⁻⁴, 0.5 × 10⁻⁴, and 10⁻⁵ gm/ml in a final volume of 2 ml of reaction mixture. Control preparations received an equal volume of acetone without DDT. Preliminary experiments revealed no enzyme inhibition with the above volume of acetone alone. Tris salt of adenosine triphosphate (ATP) (Sigma Chemicals) was used in concentrations of 2.5 mM, 1.25 mM and 0.625 mM. The phosphate released at the end of 30 minutes was determined by the method of Stanton (1968).5

III. RESULTS

The inhibitory effect of DDT on Ca ATPase is shown in figure l. When l/velocity was plotted against inhibitor concentration, the inhibitor binding constant (I) was approximately 8.8 × 10⁻⁴ gm/ml. The competitive nature of the inhibition of Ca ATPase by DDT is shown when the reciprocal of velocity is plotted versus the reciprocal of substrate concentration (figure 2). This plot shows that the inhibition by DDT was overcome by addition of more substrate (ATP).




Figure 1. In Vitro Ca ATPase Inhibition by DDT




Figure 2. Lineweaver-Burke Plot of Ca ATPase Inhibition by DDT

IV. <u>DISCUSSION</u>

The results of this study have demonstrated the inhibition of Ca ATPase by DDT in an *in vitro* microsomal test system. Furthermore, this inhibition has been established to be competitive in nature. Our results are similar to those of Miller, et al., $(1976)^3$ who showed Ca ATPase inhibition by DDE in shell gland homogenate. Previous studies have shown DDT-induced eggshell thinning to be due to decreased calcium content of eggshells (Bitman, et al., 1969). Since calcium ATPase has been shown to be associated with calcium transport in the eggshell gland, the inhibition of this enzyme *in vitro* offers a possible explanation for DDT-induced eggshell thinning.

LITERATURE CITED

- 1. Kolaja, G. J. and Hinton, D. E. Morphologic Lesions in the Eggshell Gland Accompanying DDT-Induced Eggshell Thinning. Environ. Pollut. 10 (1976) in press.
- 2. Pike, J. W. and Alvarado, R. H. Ca⁺⁺-Mg₂⁺⁺ Activated ATPase in the Shell Gland of Japanese Quail. Comp. Biochem. Physiol. 51B, pp. 119-125 (1975).
- 3. Miller, D. S., Kinter, W. B. and Peakall, D. B. Enzymatic Basis for DDT-Induced Eggshell Thinning in a Sensitive Bird. Nature 259, pp. 122-124 (1976).
- 4. Rorive, G. and Kleinzeller, A. Ca₂⁺⁺ Activated ATPase From Renal Tubular Cells. Methods of Enzymology 33, pp 303-306 (1974).
- 5. Stanton, M. B. Colorimetric Determination of Inorganic Phosphate in the Presence of Biological Material and Adenosine Triphosphate. Analyt. Biochem. 22, pp 27-34 (1968).
- 6. Bitman, S., Cecil, H. C., Harris, S. J. and Fries, G. F. DDT Induces a Decrease in Eggshell Calcium. Nature 224, pp 44-45 (1969).